

Treatment Tracker

Track your treatment each day for the next few weeks

When you keep track of your treatment, you're less likely to forget to mention something important to your doctor. Writing down when you use your inhaler also helps remind you to stay on top of your doctor's recommended COPD treatment.



Check off each time you take COMBIVENT RESPIMAT

The recommended dose is 4 inhalations a day. Not to exceed 6 inhalations in 24 hours. Follow directions from your doctor.

Do not use COMBIVENT[®] RESPIMAT[®] (ipratropium bromide and albuterol) more often than your doctor has directed. Deaths have been reported with similar inhaled medicines in asthma patients who use the medicine too much. Seek immediate medical attention if your treatment with COMBIVENT RESPIMAT becomes less effective for symptomatic relief, your symptoms become worse, and/or you need to use the product more frequently than usual.



Prepare for your next appointment with your doctor

As you record your doses each day, keep in mind anything you'd like to tell your doctor about your treatment.

In your discussion with your doctor, you may want to be prepared to answer questions such as:

- **Are you having more or less trouble breathing?**
- **Are you having trouble doing everyday activities?**
- **Are you using your rescue inhaler more or less often?**

	MONDAY	TUESDAY	WEDNESDAY	THURSDAY	FRIDAY	SATURDAY	SUNDAY	
WEEK 1	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
WEEK 2	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>

This treatment tracker has been designed based on the recommended dosing of 1 inhalation 4 times a day. You should always follow the instructions provided by your doctor.

Treatment Tracker

Track your treatment each day for the next few weeks

When you keep track of your treatment, you're less likely to forget to mention something important to your doctor. Writing down when you use your inhaler also helps remind you to stay on top of your doctor's recommended COPD treatment.



Check off each time you take COMBIVENT RESPIMAT

The recommended dose is 4 inhalations a day. Not to exceed 6 inhalations in 24 hours. Follow directions from your doctor.

Do not use COMBIVENT[®] RESPIMAT[®] (ipratropium bromide and albuterol) more often than your doctor has directed. Deaths have been reported with similar inhaled medicines in asthma patients who use the medicine too much. Seek immediate medical attention if your treatment with COMBIVENT RESPIMAT becomes less effective for symptomatic relief, your symptoms become worse, and/or you need to use the product more frequently than usual.



Prepare for your next appointment with your doctor

As you record your doses each day, keep in mind anything you'd like to tell your doctor about your treatment.

In your discussion with your doctor, you may want to be prepared to answer questions such as:

- **Are you having more or less trouble breathing?**
- **Are you having trouble doing everyday activities?**
- **Are you using your rescue inhaler more or less often?**

	MONDAY	TUESDAY	WEDNESDAY	THURSDAY	FRIDAY	SATURDAY	SUNDAY	
WEEK 3	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
WEEK 4	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	COMBIVENT RESPIMAT DOSES: <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>

This treatment tracker has been designed based on the recommended dosing of 1 inhalation 4 times a day. You should always follow the instructions provided by your doctor.

INDICATION FOR USE

COMBIVENT® RESPIMAT® (ipratropium bromide and albuterol) is indicated for use in patients with chronic obstructive pulmonary disease (COPD) on a regular aerosol bronchodilator (inhaler) who continue to have evidence of bronchospasm (airway narrowing) and who require a second bronchodilator.

IMPORTANT SAFETY INFORMATION

Do not use COMBIVENT RESPIMAT if you are allergic to any of the ingredients in COMBIVENT RESPIMAT or to atropine or other similar drugs.

COMBIVENT RESPIMAT can cause the narrowing of the airways to get worse (paradoxical bronchospasm) which may be life threatening. If this happens, stop taking COMBIVENT RESPIMAT at once and call your doctor or get emergency help.

COMBIVENT RESPIMAT can cause serious heart-related side effects, such as palpitations, chest pain, rapid heart rate, high blood pressure, tremor, or nervousness. Call your doctor if you experience any of these symptoms.

COMBIVENT RESPIMAT may increase eye pressure, which may cause or worsen some types of glaucoma.

Do not get the spray into your eyes.

COMBIVENT RESPIMAT may cause difficulty with urination.

If you have vision changes or eye pain or if you have difficulty with urination, stop taking COMBIVENT RESPIMAT and call your doctor right away.

Dizziness and blurred vision may occur with COMBIVENT RESPIMAT. Should you experience these symptoms, use caution when engaging in activities such as driving a car or operating appliances or other machines.

Do not use COMBIVENT RESPIMAT more often than your doctor has directed. Deaths have been reported with similar inhaled medicines in asthma patients who use the medicine too much. Seek immediate medical attention if your treatment with COMBIVENT RESPIMAT becomes less effective for

symptomatic relief, your symptoms become worse, and/or you need to use the product more frequently than usual.

Allergic reactions may occur, including itching, swelling of the face, lips, tongue, or throat (involving difficulty in breathing or swallowing), rash, hives, bronchospasm (airway narrowing), or anaphylaxis. Some of these may be serious. If you experience any of these symptoms, stop taking COMBIVENT RESPIMAT at once and call your doctor or get emergency help.

Tell your doctor about all your medical conditions, especially if you have narrow-angle glaucoma, prostate or urinary problems, a history of heart conditions (such as irregular heartbeat, high blood pressure), seizures, thyroid disorder, diabetes, low potassium levels, or kidney or liver disease. Also tell your doctor if you are pregnant or nursing.

Tell your doctor about all medicines you are taking, especially heart medications or drugs to treat depression.

The most common side effects reported with use of COMBIVENT RESPIMAT include infection of the ears, nose, and throat, runny nose, cough, bronchitis, headache, and shortness of breath.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call **1-800-FDA-1088**.

Read the step-by-step Instructions for Use for COMBIVENT RESPIMAT before you use your inhaler.

Please see full [Prescribing Information](#) including Instructions for Use.